

# Infinite generation of the kernels of the Magnus and Burau representations

Thomas Church and Benson Farb \*

September 26, 2009

## Abstract

Consider the kernel  $\text{Mag}_g$  of the Magnus representation of the Torelli group and the kernel  $\text{Bur}_n$  of the Burau representation of the braid group. We prove that for  $g \geq 2$  and for  $n \geq 6$  the groups  $\text{Mag}_g$  and  $\text{Bur}_n$  have infinite rank first homology. As a consequence we conclude that neither group has any finite generating set. The method of proof in each case consists of producing a kind of “Johnson-type” homomorphism to an infinite rank abelian group, and proving the image has infinite rank. For the case of  $\text{Bur}_n$ , we do this with the assistance of a computer calculation.

## 1 Introduction

**The Magnus kernel.** Let  $S := S_{g,1}$  be a compact, connected, oriented surface of genus  $g \geq 2$  with one boundary component. Let  $\text{Mod}_{g,1}$  denote the *mapping class group* of  $S$ , which is the group of homotopy classes of orientation-preserving homeomorphisms of  $S$  which fix  $\partial S$  pointwise. Let  $\mathcal{I}_{g,1}$  denote the *Torelli group*, which is the subgroup of  $\text{Mod}_{g,1}$  consisting of elements that act trivially on  $H := H_1(S, \mathbb{Z})$ .

$\text{Mod}_{g,1}$  acts on the fundamental group  $\pi_1(S)$ , inducing an action on the solvable quotient  $\Gamma/\Gamma^3$ , where  $\Gamma := \pi_1(S)$ ,  $\Gamma^2 = [\Gamma, \Gamma]$  and  $\Gamma^3 = [\Gamma^2, \Gamma^2]$  are the first three terms of the derived series of  $\Gamma$ . In this paper we consider the group

$$\text{Mag}_g := \text{kernel}(\text{Mod}(S) \rightarrow \text{Aut}(\Gamma/\Gamma^3)).$$

In 1939, Magnus ([Ma]; see also [Bi, Chapter 3]) used the Fox calculus to construct a representation

$$r: \mathcal{I}_{g,1} \rightarrow \text{GL}_{2g}(\mathbb{Z}H)$$

now called the *Magnus representation*. It follows from [Fox, Theorem 4.9] that the kernel of  $r$  coincides with  $\text{Mag}_g$ . This group is called the *Magnus kernel*.

It was an open question for some time whether or not  $\text{Mag}_g$  is nontrivial. This was settled in the affirmative by Suzuki in [S1]. The first main result of this paper is that  $\text{Mag}_g$  is in fact quite large.

---

\*The second author gratefully acknowledges support from the National Science Foundation.

**Theorem 1.1.** *For  $g \geq 2$  the group  $H_1(\text{Mag}_g, \mathbb{Z})$  has infinite rank.*

As the abelianization of a finitely-generated group has finite rank, we deduce the following.

**Corollary 1.2.** *For  $g \geq 2$  the group  $\text{Mag}_g$  has no finite generating set.*

The idea of our proof of Theorem 1.1 is to define a kind of “Johnson-type” homomorphism (see [J1]):

$$\Psi: \text{Mag}_g \rightarrow \text{Hom}(G^{\text{ab}}, \bigwedge^2 G^{\text{ab}})$$

where  $G = [\Gamma, \Gamma]$  and  $G^{\text{ab}}$  denotes the abelianization of  $G$ . We then construct infinitely many linearly independent elements contained in the image.

**The Burau kernel.** Let  $B_n$  denote the braid group on  $n$  strands.  $B_n$  can be realized (see Section 4 below) as a subgroup of the automorphism group  $\text{Aut}(F_n)$  of the free group of rank  $n$ . The *Burau representation* is a homomorphism

$$\rho_n: B_n \rightarrow \text{GL}_n(\mathbb{Z}[t, t^{-1}]).$$

We define the *Burau kernel*, denoted  $\text{Bur}_n$ , to be the kernel of  $\rho_n$ . Let  $K$  be the kernel of the homomorphism  $F_n \rightarrow \mathbb{Z}$  taking each fixed generator of  $F_n$  to 1. It follows easily from [Fox] that

$$\text{Bur}_n = \text{kernel}(B_n \rightarrow \text{Aut}(F_n/[K, K])).$$

While  $\rho_3$  is faithful, it was a longstanding problem as to whether or not  $\rho_n$  is faithful (i.e. whether  $\text{Bur}_n$  is nontrivial) for  $n > 3$ . This was solved by Moody [Mo], Long–Paton [LP], and Bigelow [Big] in various cases, with the result that  $\text{Bur}_n$  is nontrivial for  $n \geq 5$ ; the case of  $n = 4$  is still open. Our next main result is that  $\text{Bur}_n$  is in fact quite large for  $n \geq 6$ .

**Theorem 1.3.** *For  $n \geq 6$  the group  $H_1(\text{Bur}_n, \mathbb{Z})$  has infinite rank; in particular,  $\text{Bur}_n$  has no finite generating set.*

To prove Theorem 1.3 we construct, similarly to the proof of Theorem 1.1 above, a homomorphism

$$\Phi: \text{Bur}_n \rightarrow \text{Hom}(K^{\text{ab}}, \bigwedge^2 K^{\text{ab}}).$$

The elements which have been constructed in the kernel of the Burau representation are geometrically elegant, but algebraically very complicated; for example, the element of  $\text{Bur}_7$  found by Long–Paton can be described by a single diagram, but as a free group automorphism sends generators of  $F_7$  to words of length up to 475137. Thus we need the assistance of a computer in order to calculate  $\Phi$  explicitly (see Section 4 below for a full discussion). For the computations in this paper we use a simpler element  $\phi_B \in \text{Bur}_n$  for  $n \geq 6$  found by Bigelow, which takes generators to words of length no more than 9841. Once we compute the form of  $\Phi(\phi_B)$ , we then use an equivariance property of  $\Phi$  to prove that the image of  $\Phi$  has infinite rank, from which Theorem 1.3 follows.

We remark that, as Problem 6.24 of [Mor], Morita posed the problem of determining the kernel of the Magnus and Burau (among other) representations. Theorem 1.1 and Theorem 1.3 can be viewed as a partial answer to this problem.

**Acknowledgements.** We are grateful to William Goldman, whose Mathematica notebook `FreeGroupAutos.nb` was very helpful in our computations of the expression in Appendix A. We would also like to thank Dan Margalit and Tam Nguyen Phan for careful comments on an earlier version of this paper.

## 2 Defining the homomorphisms

The following construction works whenever one considers a group of automorphisms of the universal 2-step nilpotent quotient of a group  $G$  acting trivially on its abelianization. Johnson [J1] considered the case  $G = \Gamma = \pi_1(S)$ .

With  $\Gamma$  equal to  $\pi_1(S)$  or  $F_n$  as in the introduction, we take  $G := [\Gamma, \Gamma]$  or  $G := K$  respectively. In either case, let  $G_i$  be the lower central series of  $G$ , defined inductively by  $G_1 = G$  and  $G_{i+1} = [G, G_i]$ . Consider the exact sequence

$$1 \rightarrow G_2 \rightarrow G \rightarrow G^{\text{ab}} \rightarrow 1. \quad (1)$$

Centralizing (1) gives

$$1 \rightarrow G_2/G_3 \rightarrow G/G_3 \rightarrow G^{\text{ab}} \rightarrow 1. \quad (2)$$

Since  $G$  is free, taking (1) as a presentation for  $G^{\text{ab}}$ , Hopf's formula gives that

$$G_2/G_3 \approx \bigwedge^2 G^{\text{ab}}.$$

$\text{Aut}(\Gamma)$  acts on  $\Gamma$ , and thus on  $G$ , and the isomorphism  $\nu: G_2/G_3 \approx \bigwedge^2 G^{\text{ab}}$  respects the action of  $\text{Aut}(\Gamma)$  on both sides. In particular, conjugation by  $\Gamma$  descends to an action on  $G^{\text{ab}}$  by  $H = \Gamma/[\Gamma, \Gamma]$  or by  $\mathbb{Z} = \Gamma/K$  respectively. In the case  $G = [\Gamma, \Gamma]$ , the fact that  $\text{Mag}_g$  acts trivially on  $\Gamma/\Gamma^3$  implies that  $\text{Mag}_g$  acts trivially on  $G^{\text{ab}} = \Gamma^2/\Gamma^3$  and on  $\bigwedge^2 G^{\text{ab}}$ . Similarly, in the case  $G = K$ , we have that  $\text{Bur}_n$  acts trivially on  $G^{\text{ab}}$  and on  $\bigwedge^2 G^{\text{ab}}$ .

Let  $f \in \text{Mag}_g$  (resp.  $f \in \text{Bur}_n$ ) be given. For  $x \in G^{\text{ab}}$ , pick any lift  $\tilde{x} \in G$ . Since  $f$  acts trivially on both the quotient and kernel of (2), we see that  $f(\tilde{x})\tilde{x}^{-1}$  lies in the kernel  $G_2/G_3$ , which we identify with  $\bigwedge^2 G^{\text{ab}}$  via the isomorphism above. One checks, exactly as in [J1], that

$$\delta_f: G^{\text{ab}} \rightarrow \bigwedge^2 G^{\text{ab}}$$

defined by  $\delta_f(x) := f(\tilde{x})\tilde{x}^{-1}$  is a well-defined homomorphism; in fact, the resulting map  $\delta_f$  is  $\mathbb{Z}H$ -linear (resp.  $\mathbb{Z}[t, t^{-1}]$ -linear) with respect to the conjugation action on  $G^{\text{ab}}$ . This is equivalent to the claim that

$$\delta_f(\gamma x \gamma^{-1}) \equiv \gamma \delta_f(x) \gamma^{-1} \bmod G_3,$$

which can be checked as follows. The difference between the left and right side is

$$(f(\gamma x \gamma^{-1}) \gamma x^{-1} \gamma^{-1}) (\gamma f(x) x^{-1} \gamma^{-1})^{-1} = f(\gamma) f(x) f(\gamma)^{-1} \gamma f(x)^{-1} \gamma^{-1},$$

which is conjugate to  $[\gamma^{-1} f(\gamma), f(x)]$ . The condition on  $f$  implies that  $f(\gamma) \equiv \gamma \bmod G_2$ , so  $\gamma^{-1} f(\gamma) \in G_2$  and  $[\gamma^{-1} f(\gamma), f(x)] \in G_3$  as desired.

One also checks, exactly as in [J1], that in the case  $G = [\Gamma, \Gamma]$ , defining the map  $\Psi$  by  $\Psi(f) := \delta_f$  gives a well-defined homomorphism;

$$\Psi: \text{Mag}_g \rightarrow \text{Hom}(G^{\text{ab}}, \bigwedge^2 G^{\text{ab}}). \quad (3)$$

and, in the case  $G = K$ , defining  $\Phi(g) := \delta_f$  gives a well-defined homomorphism:

$$\Phi: \text{Bur}_n \rightarrow \text{Hom}(G^{\text{ab}}, \bigwedge^2 G^{\text{ab}}). \quad (4)$$

The homomorphisms  $\Psi$  and  $\Phi$  are equivariant with respect to the natural  $\text{Aut}(\Gamma)$ -actions on the source and target.

### 3 Computing the image of $\Psi$

Let  $S_{0,4}$  denote the 2-sphere with 4 open disks removed. A *lantern* in  $S$  is an embedding  $S_{0,4} \hookrightarrow S$ . Consider the two simple closed curves  $\alpha$  and  $\beta$  and the three arcs  $A_1, A_2$  and  $A_3$  on  $S_{0,4}$  given in Figure 1.




Figure 1: The simple closed curves  $\alpha$  and  $\beta$ , and the arcs  $A_1, A_2, A_3$ .

One directly computes the action of  $f := T_\alpha T_\beta^{-1}$  on  $A_1, A_2$  and  $A_3$ , as follows (see Figure 2). Let  $x, y$ , and  $z$  be the loops which begin with  $A_1, A_2$  and  $A_3$ , respectively, go clockwise around the appropriate boundary component of  $S_{0,4}$ , then come back along the same arc  $A_i$ . Let  $X, Y, Z$  be the inverses of  $x, y, z$  in  $\pi_1(S_{0,4})$ . Then:

$$f(A_1) = xyXzxYXZA_1 = [xyX, z]A_1$$

$$f(A_2) = ZXzxA_2 = [Z, X]A_2$$

$$f(A_3) = ZXzxYXZxzyXA_3 = [ZXz, xYX]A_3$$

Let  $L$  be an embedding of a lantern in  $S$  with the property that each of the four boundary curves of  $L$  are separating in  $S$ .<sup>1</sup> In this case we can observe that  $T_\alpha T_\beta^{-1} \in \text{Mag}_g$ , as follows.

<sup>1</sup>To formally identify  $x, y, z$  with elements of  $\Gamma = \pi_1(S)$ , we choose a basepoint on  $\partial S$ , and arcs from this basepoint to  $L$  meeting  $L$  in one point. Since  $f$  is the identity off of  $L$ , any ambiguity in the choice of these paths to  $L$  does not affect the computation.




Figure 2: The arcs  $f(A_1)$ ,  $f(A_2)$  and  $f(A_3)$ .

Note that the elements corresponding to  $x, y, z$  all lie in  $\Gamma^2$ . Furthermore,  $\Gamma = \pi_1(S)$  has a basis where each element  $c$  is either disjoint from  $L$ , or else of the form  $c = A\gamma A^{-1}$ , where  $A$  is an arc intersecting  $L$  in some  $A_i$  and  $\gamma$  is a loop disjoint from  $L$ . In the former case the element  $f = T_\alpha T_\beta^{-1}$  fixes  $c$ . In the latter case, assume for example that  $A$  intersects  $L$  in  $A_2$ ; then we have

$$f(c) = f(A\gamma A^{-1}) = f(A)\gamma f(A)^{-1} = [Z, X]A\gamma A^{-1}[X, Z] = [Z, X]c[X, Z]$$

Since  $x, y, z \in \Gamma^2$ , we have  $[Z, X] \in \Gamma^3$ ; thus  $f(c) \equiv c \pmod{\Gamma^3}$ . The same is true for  $A_1$  and  $A_3$ , so we conclude that  $f(c) \equiv c \pmod{\Gamma^3}$  for all elements of a basis for  $\Gamma$ , implying  $T_\alpha T_\beta^{-1} \in \text{Mag}_g$ . Suzuki gave a more illuminating proof that elements of this form lie in  $\text{Mag}_g$  in [S2].

We are now ready to compute  $\Psi$ . For  $a, b \in \Gamma$ , we denote by  $\{a, b\}$  the image of  $[a, b] \in G$  in  $G^{\text{ab}}$  under the abelianization map.

**Proposition 3.1.** *Let  $L$  be a lantern embedded in  $S$  so that each of the four boundary curves of  $L$  are separating in  $S$ . Let  $a$  and  $b$  be loops intersecting  $L$  in  $A_1$  and  $A_2$ . Then*

$$\Psi(T_\alpha T_\beta^{-1})(\{a, b\}) = (a - 1)(b - 1)[x \wedge z + y \wedge z] \quad (5)$$

Note that the right hand side of (5) is an element of  $\bigwedge^2 G^{\text{ab}}$ , considered as a  $\mathbb{Z}H$ -module, and  $a, b$  are taken to be elements of  $H$ .

*Proof.* As in the computation above, we have

$$f([a, b]) = [f(a), f(b)] = [wa, vb]$$

where

$$w = [[xyX, z], a] \quad \text{and} \quad v = [[Z, X], b].$$

From the assumption on the embedding of  $L$  we have  $x, y, z \in G$ , and thus  $w, v \in G_2$ . We will use the following commutator identities, which hold in any group; we write  ${}^x y$  for  $xyx^{-1}$ .

$$[wa, b] = {}^w[a, b] [w, b] \quad [a, vb] = [a, v] {}^v[a, b]$$




Figure 3: The curves  $\gamma$  and  $\delta_k$  for  $k = 3$ .

We then find that

$$[wa, vb] = {}^w[a, v] {}^{wv}[a, b] [w, v] {}^v[w, b]$$

Note that the second term lies in  $G$ , the first and fourth in  $G_2$ , and the third in  $G_3$ .

We want to compute  $f([a, b])[a, b]^{-1}$  as an element of  $G_2/G_3$ . Note that  $[w, v] \equiv 0 \pmod{G_3}$ , and that conjugating an element of  $G$  by an element of  $G_2$  is a trivial operation modulo  $G_3$ . Finally, since  $[[a, b], [w, b]] \in G_3$ , we can move  $[a, b]$  to the right to cancel  $[a, b]^{-1}$ . We thus obtain

$$\begin{aligned} f([a, b])[a, b]^{-1} &= {}^w[a, v] {}^{wv}[a, b] [w, v] {}^v[w, b] [a, b]^{-1} \\ &\equiv [a, v][a, b][w, b][a, b]^{-1} \pmod{G_3} \\ &\equiv [a, v][w, b] \pmod{G_3}. \end{aligned}$$

Recall that action of  $\Gamma$  on  $\Gamma$  by conjugation descends to a  $\mathbb{Z}H$  action on  $G^{\text{ab}}$ . Recall from above the isomorphism  $\nu: G_2/G_3 \rightarrow \bigwedge^2 G^{\text{ab}}$ . Since the homology class of  $x$  is trivial in  $H$ , we have

$$\nu([xyX, z]) = y \wedge z \quad \text{and} \quad \nu([Z, X]) = z \wedge x.$$

It follows that

$$\nu(w) = \nu([xyX, z], a) = (1 - a)y \wedge z$$




Figure 4: The boundary curves of  $L_k$ ; the subsurfaces cut off by these curves are shaded.

and

$$\nu(v) = \nu([Z, X], b) = (1 - b)z \wedge x.$$

We therefore have that

$$\nu([a, v][w, b]) = (a - 1)v - (b - 1)w = (a - 1)(1 - b)z \wedge x - (b - 1)(1 - a)y \wedge z.$$

We conclude that

$$\Psi(T_\alpha T_\beta^{-1})(\{a, b\}) = (a - 1)(b - 1)[x \wedge z + y \wedge z]$$

as desired.  $\square$

**Theorem 3.2.** *The image of  $\Psi$  has infinite rank for  $g \geq 3$ .*

*Proof.* Let  $\gamma$  and  $\delta_k$  be the curves depicted in Figure 3. The figure depicts the case  $k = 3$ ; in general  $\delta_k$  has  $k$  twists around the upper right handle. (Specifically, the curve  $\delta_k$  is equal to  $T_{a_3}^k(\delta_0)$ , where  $a_3$  is as in Figure 5.) The regular neighborhood of  $\gamma \cup \delta_k$  is a lantern  $L_k$ , and we fix an identification of  $L_k$  with our reference lantern  $L$  by specifying that  $\gamma$  and  $\delta_k$  should




Figure 5: A basis for  $\pi_1(S_{g,1})$ .

correspond to  $xy$  and  $yz$  respectively. Let  $f_k \in \text{Mag}_g$  be the element corresponding under this identification to the mapping class  $T_\alpha T_\beta^{-1}$  on  $L$ ; it is easy to check using the lantern relation that  $f_k$  is in fact  $[T_\gamma^{-1}, T_{\delta_k}^{-1}]$ . We will show that the images  $\Psi(f_k)$  are linearly independent (over  $\mathbb{Z}$ ).

The boundary curves of  $L_k$  are depicted in Figure 4. With the basis  $a_1, b_1, \dots, a_g, b_g$  for  $\pi_1(S_{g,1})$  as illustrated in Figure 5, we see that as curves  $x, y$  and  $z$  can be represented by  $[a_1, b_1]$ ,  $[a_2, b_3 a_3^k b_2]$ , and  $[b_2 a_2^{-1} b_2^{-1} a_3, b_3 a_3^k]$  respectively. As based loops, we actually have the conjugate  $z = {}^c[b_2 a_2^{-1} b_2^{-1} a_3, b_3 a_3^k]$ , where  $c = [b_3, a_3][b_2, a_2]a_2$ . Note that with this representative for  $z$ , we have  $xyz = [a_1, b_1][a_2, b_2][a_3, b_3]$ , the fourth boundary curve in Figure 4.

Note that  $a_1$  and  $a_2$  intersect each  $L_k$  in arcs corresponding to  $A_1$  and  $A_2$ . Thus by Proposition 3.1, we have that

$$\Psi(f_k)(\{a_1, a_2\}) = (a_1 - 1)(a_2 - 1)[(\{a_1, b_1\} + \{a_2, b_3 a_3^k b_2\}) \wedge a_2 \{b_2 a_2^{-1} b_2^{-1} a_3, b_3 a_3^k\}]$$

Denote this element of  $\bigwedge^2 G^{\text{ab}}$  by  $\alpha_k$ . We now check that  $\{\alpha_k\}$  is linearly independent as follows. There is a standard embedding  $G^{\text{ab}} \hookrightarrow (\mathbb{Z}H)^{2g}$  given by sending the class  $[x]$  to  $(\partial x / \partial z_1, \dots, \partial x / \partial z_n)$ , where  $\{z_i\}$  is our basis for  $F_n$  and where  $\partial / \partial z_i$  are the Fox derivatives (see e.g. [CP] for a detailed explanation of this embedding). The only property of this embedding that we will need is that the components that make up  $\alpha_k$  are mapped as follows by the embedding. Here the  $A_i$  and  $B_i$  make up a basis for  $(\mathbb{Z}H)^{2g}$ .

$$\begin{aligned} \{a_1, b_1\} &\mapsto (1 - b_1)A_1 - (1 - a_1)B_1 \\ \{a_2, b_3 a_3^k b_2\} &\mapsto (1 - b_3 a_3^k b_2)A_2 \\ &\quad - (1 - a_2)(B_3 + b_3(1 + \dots + a_3^{k-1})A_3 + b_3 a_3^k B_2) \\ \{b_2 a_2^{-1} b_2^{-1} a_3, b_3 a_3^k\} &\mapsto (1 - b_3 a_3^k)((1 - a_2^{-1})B_2 - a_2^{-1} b_2 A_2 + a_2^{-1} A_3) \\ &\quad - (1 - a_2^{-1} a_3)(B_3 + b_3(1 + \dots + a_3^{k-1})A_3) \end{aligned}$$






Figure 6: The commutator  $[T_\gamma, T_\delta]$  lies in  $\text{Mag}_2$ .

By expanding out  $\alpha_k$ , we see that  $\alpha_N$  is the only such element which contains the term  $A_1 \wedge b_3 a_3^N B_2$  with nonzero coefficient; it follows that the  $\alpha_k$  are linearly independent, as desired.  $\square$

As the image of  $\Psi$  is abelian, Theorem 3.2 immediately implies Theorem 1.1 for  $g \geq 3$ . Note that the proof of Theorem 3.2 used in an essential way that  $g \geq 3$ . So in order to complete the proof of Theorem 1.1, we need another argument when  $g = 2$ .

**Theorem 3.3.**  $H_1(\text{Mag}_2)$  has infinite rank.

*Proof.* Suzuki showed that the element  $f = [T_\gamma, T_\delta]$  is in  $\text{Mag}_2$  for  $\gamma$  and  $\delta$  as in Figure 6; in particular  $\text{Mag}_2$  is nontrivial. Let  $S_2$  be a closed surface of genus 2; we denote by  $\mathcal{I}_{2,*}$  the Torelli group of  $S_2$  with respect to a marked point  $*$ , and by  $\mathcal{I}_2$  the Torelli group of the closed surface  $S_2$ . By Johnson [J2], we have the exact sequence

$$1 \rightarrow \mathbb{Z} \rightarrow \mathcal{I}_{2,1} \xrightarrow{p} \mathcal{I}_{2,*} \rightarrow 1,$$

where the kernel is generated by a twist  $T_\omega$  around the boundary  $\omega = \partial S_2$ . It is easy to check that the action of  $T_\omega$  on  $\pi_1(S_{2,1})$  is conjugation by  $\omega$ ; since  $\omega \notin \Gamma^3$ , we see that  $T_\omega \notin \text{Mag}_2$ . It follows that  $p$  restricts to an isomorphism between  $\text{Mag}_2$  and a subgroup  $p(\text{Mag}_2) < \mathcal{I}_{2,*}$ .

Again by Johnson [J2], we have the exact sequence

$$1 \rightarrow \Lambda \rightarrow \mathcal{I}_{2,*} \xrightarrow{\pi} \mathcal{I}_2 \rightarrow 1,$$

where  $\Lambda \approx \pi_1(S_2, *)$ ; note that  $\mathcal{I}_{2,*}$  acts on  $\pi_1(S_2, *)$ , and the restriction to  $\Lambda$  is just the action by conjugation. Mess [Me] proved that  $\mathcal{I}_2$  is free of infinite rank. It is easy to see from Figure 6 that  $f \in \ker \pi = \Lambda$ . We use the following well-known lemma.

**Lemma 3.4.** *Any nontrivial infinite index normal subgroup of a surface group or free group is an infinite rank free group.*

If  $\pi \circ p(\text{Mag}_2) < \mathcal{I}_2 \approx F_\infty$  is nontrivial, then by Lemma 3.4,  $\text{Mag}_2$  surjects to the infinite rank free group  $\pi \circ p(\text{Mag}_2)$ , and we are done.

Suppose that  $p(\text{Mag}_2) \subset \ker \pi = \Lambda$ . Any  $\varphi \in \text{Mag}_2$  acts trivially on  $\Gamma/\Gamma^3$ ; thus  $p(\varphi)$  acts trivially on  $\pi_1(S_2)/\pi_1(S_2)^3$ . Since the action of  $\Lambda$  is by conjugation, this implies that  $p(\varphi)$  lies in  $\Lambda^3$ . Thus  $p(\text{Mag}_2)$  has infinite index in  $\Lambda$ , and so by Lemma 3.4,  $p(\text{Mag}_2) \approx \text{Mag}_2$  is an infinite rank free group.  $\square$

Theorem 1.1, and hence Corollary 1.2, follows immediately from Theorems 3.2 and 3.3.

**Remark.** One can check by explicit computation that for Suzuki's element  $f \in \text{Mag}_2$  above,  $\Psi(f) = 0$ . It would be interesting to know whether  $\Psi$  in fact vanishes on  $\text{Mag}_2$ .

## 4 Computing the image of $\Phi$

The kernel  $K$  of the map from  $F_n = \langle x_1, \dots, x_n \rangle$  to  $\mathbb{Z} = \langle t \rangle$  which sends each  $x_i \mapsto t$  is normally generated by the elements  $x_i x_j^{-1}$ . If we set  $x_{i,k} := x_1^k x_i x_1^{-k-1}$  for  $i \neq 1$  and  $k \in \mathbb{Z}$ , then  $\{x_{i,k}\}$  gives a basis for  $K$  as a free group. As above, the conjugation of  $K$  by  $F_n$  descends to a  $\mathbb{Z}[t, t^{-1}]$  action on  $K^{\text{ab}}$ . With respect to this action we have  $x_{i,k} = t^k x_{i,0}$ , and thus  $K^{\text{ab}}$  is a free  $\mathbb{Z}[t, t^{-1}]$ -module with basis  $\{y_i = x_{i,0}\}_{i \neq 1}$ .

The braid group  $B_n$  has generators  $\sigma_1, \dots, \sigma_{n-1}$ ; the action of  $\sigma_i$  on  $F_n$  sends  $x_i \mapsto x_i x_{i+1} x_i^{-1}$ ,  $x_{i+1} \mapsto x_i$ , and fixes the other generators. The action of  $B_n$  on  $K^{\text{ab}}$  commutes with the  $\mathbb{Z}[t, t^{-1}]$  action.




Figure 7: The two arcs defining Bigelow's element  $\phi_B$ .

**Theorem 4.1.** *The image of  $\Phi$  has infinite rank for  $n \geq 6$ .*

*Proof.* The element of the kernel found by Bigelow in [Big] is the commutator of the half-twists along the arcs displayed in Figure 7. In terms of the Artin generators, this is

$$\phi_B = [\psi_1 \sigma_3^{-1} \psi_1^{-1}, \psi_2 \sigma_3^{-1} \psi_2^{-1}], \quad \text{where } \psi_1 = \sigma_4 \sigma_5^{-1} \sigma_2^{-1} \sigma_1 \quad \text{and } \psi_2 = \sigma_4^{-1} \sigma_5^2 \sigma_2 \sigma_1^{-1}.$$

In Appendix A, we give the computation of  $\alpha := \Phi(\phi_B)([x_2x_1^{-1}]) = \Phi(\phi_B)(y_2)$ ; it has 262 terms. The only fact about  $\alpha$  that we will need is that its highest term of the form  $y_2 \wedge t^k y_4$  is  $-2y_2 \wedge t^3 y_4$ , and its highest term of the form  $y_2 \wedge t^k y_5$  is  $+2y_2 \wedge t^2 y_5$  (these terms are set in boxes in the appendix).

It is easy to check that

$$\sigma_4^2(x_4) = x_4 x_5 x_4 x_5^{-1} x_4^{-1}$$

$$\sigma_4^2(x_5) = x_5 x_4 x_5^{-1}$$

$$\sigma_4^2(x_i) = x_i \text{ for } i \neq 4, 5.$$

By induction, for  $k \geq 1$  we have

$$\sigma_4^{2k}(x_4) = (x_4 x_5)^k x_4 (x_4 x_5)^{-k}$$

$$\sigma_4^{2k}(x_5) = (x_4 x_5)^{k-1} x_4 x_5 x_4^{-1} (x_4 x_5)^{k-1}$$

$$\sigma_4^{2k}(x_i) = x_i \text{ for } i \neq 4, 5.$$

The action of  $\sigma_4^{2k}$  on  $K^{\text{ab}}$  in terms of our basis is thus given by:

$$\begin{aligned} y_4 &\mapsto (1 - t + t^2 - \dots - t^{k-1} + t^k) y_4 &+& (t - t^2 + \dots + t^{k-1} - t^k) y_5 \\ y_5 &\mapsto (1 - t + t^2 - \dots - t^{k-1}) y_4 &+& (t - t^2 + \dots + t^{k-1}) y_5 \\ y_i &\mapsto y_i \text{ for } i \neq 4, 5 \end{aligned}$$

Now for  $k \geq 0$  set

$$\alpha_k := \Phi(\sigma_4^{2k} \phi_B \sigma_4^{-2k})(y_2).$$

By the equivariance of  $\Phi$ , and since  $\sigma_4$  fixes  $y_2$ , we have  $\alpha_k = \sigma_4^{2k} \cdot \alpha$ . From the action of  $\sigma_4^{2k}$  on  $K^{\text{ab}}$ , we can see that the highest term in  $\alpha_N$  of the form  $y_2 \wedge t^k y_4$  will be  $-2y_2 \wedge t^{3+N} y_4$ . Thus  $\alpha_N$  is not contained in the span of  $\{\alpha_1, \dots, \alpha_{N-1}\}$ ; it follows that the  $\alpha_k$  are linearly independent over  $\mathbb{Z}$ , and thus the image of  $\Phi$  has infinite rank.  $\square$

Theorem 1.3 follows immediately.

## A Appendix

The following computation was made, with the method explained in Section 4, with the help of *Mathematica*. A *Mathematica* notebook implementing these computations can be found at:

<http://math.uchicago.edu/~tchurch/infinitegeneration.html>

The output of this notebook is  $\Phi(\phi_B)(y_2)$ , which is:

$$\begin{array}{cccccc} -t^{-3}y_2 \wedge t^{-2}y_2 & +t^{-3}y_2 \wedge t^{-1}y_2 & -t^{-3}y_2 \wedge y_2 & -t^{-2}y_2 \wedge y_2 & +t^{-1}y_2 \wedge y_2 & \\ +t^{-2}y_2 \wedge ty_2 & +t^{-1}y_2 \wedge ty_2 & -2y_2 \wedge t^2y_2 & +ty_2 \wedge t^3y_2 & +t^2y_2 \wedge t^3y_2 & \\ -t^3y_2 \wedge t^4y_2 & +t^{-3}y_2 \wedge t^{-4}y_3 & -t^{-2}y_2 \wedge t^{-4}y_3 & -t^{-3}y_2 \wedge t^{-3}y_3 & +t^{-1}y_2 \wedge t^{-3}y_3 & \end{array}$$

$+t^{-2}y_2 \wedge t^{-2}y_3$	$-t^{-1}y_2 \wedge t^{-2}y_3$	$+t^{-3}y_2 \wedge t^{-1}y_3$	$-y_2 \wedge t^{-1}y_3$	$+ty_2 \wedge t^{-1}y_3$
$-t^2y_2 \wedge t^{-1}y_3$	$-2t^{-2}y_2 \wedge y_3$	$+t^3y_2 \wedge y_3$	$+t^{-1}y_3 \wedge y_3$	$+2t^{-1}y_2 \wedge ty_3$
$-t^{-1}y_3 \wedge ty_3$	$-2y_2 \wedge t^2y_3$	$-t^4y_2 \wedge t^2y_3$	$+t^{-1}y_3 \wedge t^2y_3$	$+ty_2 \wedge t^3y_3$
$+t^4y_2 \wedge t^3y_3$	$-y_3 \wedge t^3y_3$	$+ty_3 \wedge t^3y_3$	$-t^2y_3 \wedge t^3y_3$	$+t^{-3}y_2 \wedge t^{-3}y_4$
$-t^{-2}y_2 \wedge t^{-3}y_4$	$-t^{-3}y_2 \wedge t^{-2}y_4$	$+t^{-1}y_2 \wedge t^{-2}y_4$	$+t^{-2}y_2 \wedge t^{-1}y_4$	$-t^{-1}y_2 \wedge t^{-1}y_4$
$+t^{-3}y_2 \wedge y_4$	$-y_2 \wedge y_4$	$+ty_2 \wedge y_4$	$-t^2y_2 \wedge y_4$	$-y_3 \wedge y_4$
$+ty_3 \wedge y_4$	$-t^2y_3 \wedge y_4$	$-2t^{-2}y_2 \wedge ty_4$	$+t^3y_2 \wedge ty_4$	$+t^{-1}y_3 \wedge ty_4$
$+t^3y_3 \wedge ty_4$	$+y_4 \wedge ty_4$	$+2t^{-1}y_2 \wedge t^2y_4$	$-t^{-1}y_3 \wedge t^2y_4$	$-t^3y_3 \wedge t^2y_4$
$-y_4 \wedge t^2y_4$	$-2y_2 \wedge t^3y_4$	$-t^4y_2 \wedge t^3y_4$	$+t^{-1}y_3 \wedge t^3y_4$	$+t^3y_3 \wedge t^3y_4$
$+y_4 \wedge t^3y_4$	$+ty_2 \wedge t^4y_4$	$+t^4y_2 \wedge t^4y_4$	$-y_3 \wedge t^4y_4$	$+ty_3 \wedge t^4y_4$
$-t^2y_3 \wedge t^4y_4$	$-ty_4 \wedge t^4y_4$	$+t^2y_4 \wedge t^4y_4$	$-t^3y_4 \wedge t^4y_4$	$+t^{-3}y_2 \wedge t^{-3}y_5$
$-t^{-2}y_2 \wedge t^{-3}y_5$	$+t^{-3}y_2 \wedge t^{-2}y_5$	$-t^{-2}y_2 \wedge t^{-2}y_5$	$+y_2 \wedge t^{-2}y_5$	$-t^{-4}y_3 \wedge t^{-2}y_5$
$+t^{-3}y_3 \wedge t^{-2}y_5$	$-t^{-1}y_3 \wedge t^{-2}y_5$	$-t^{-3}y_4 \wedge t^{-2}y_5$	$+t^{-2}y_4 \wedge t^{-2}y_5$	$-y_4 \wedge t^{-2}y_5$
$-t^{-3}y_5 \wedge t^{-2}y_5$	$-2t^{-3}y_2 \wedge t^{-1}y_5$	$+t^{-1}y_2 \wedge t^{-1}y_5$	$+y_2 \wedge t^{-1}y_5$	$-ty_2 \wedge t^{-1}y_5$
$+t^{-4}y_3 \wedge t^{-1}y_5$	$-t^{-2}y_3 \wedge t^{-1}y_5$	$+2y_3 \wedge t^{-1}y_5$	$+t^{-3}y_4 \wedge t^{-1}y_5$	$-t^{-1}y_4 \wedge t^{-1}y_5$
$+2ty_4 \wedge t^{-1}y_5$	$+t^{-3}y_5 \wedge t^{-1}y_5$	$+t^{-3}y_2 \wedge y_5$	$+2t^{-2}y_2 \wedge y_5$	$-2t^{-1}y_2 \wedge y_5$
$-y_2 \wedge y_5$	$-t^2y_2 \wedge y_5$	$-t^{-3}y_3 \wedge y_5$	$+t^{-2}y_3 \wedge y_5$	$-y_3 \wedge y_5$
$-ty_3 \wedge y_5$	$-t^2y_3 \wedge y_5$	$-t^{-2}y_4 \wedge y_5$	$+t^{-1}y_4 \wedge y_5$	$-ty_4 \wedge y_5$
$-t^2y_4 \wedge y_5$	$-t^3y_4 \wedge y_5$	$+t^{-1}y_5 \wedge y_5$	$-t^{-3}y_2 \wedge ty_5$	$-t^{-1}y_2 \wedge ty_5$
$+y_2 \wedge ty_5$	$+ty_2 \wedge ty_5$	$+t^3y_2 \wedge ty_5$	$+t^{-1}y_3 \wedge ty_5$	$-y_3 \wedge ty_5$
$+2ty_3 \wedge ty_5$	$+t^3y_3 \wedge ty_5$	$+y_4 \wedge ty_5$	$-ty_4 \wedge ty_5$	$+2t^2y_4 \wedge ty_5$
$+t^4y_4 \wedge ty_5$	$-t^{-2}y_5 \wedge ty_5$	$-y_5 \wedge ty_5$	$+t^{-2}y_2 \wedge t^2y_5$	$-t^{-1}y_2 \wedge t^2y_5$
$+2y_2 \wedge t^2y_5$	$-t^2y_2 \wedge t^2y_5$	$+t^3y_2 \wedge t^2y_5$	$-t^{-1}y_3 \wedge t^2y_5$	$-t^2y_3 \wedge t^2y_5$
$-y_4 \wedge t^2y_5$	$-t^3y_4 \wedge t^2y_5$	$+t^{-1}y_5 \wedge t^2y_5$	$-2y_5 \wedge t^2y_5$	$+ty_5 \wedge t^2y_5$
$-ty_2 \wedge t^3y_5$	$-t^2y_2 \wedge t^3y_5$	$-t^4y_2 \wedge t^3y_5$	$+y_3 \wedge t^3y_5$	$+ty_4 \wedge t^3y_5$
$+ty_5 \wedge t^3y_5$	$+t^2y_5 \wedge t^3y_5$	$+t^2y_2 \wedge t^4y_5$	$+t^3y_2 \wedge t^4y_5$	$-ty_3 \wedge t^4y_5$
$+t^3y_3 \wedge t^4y_5$	$-t^2y_4 \wedge t^4y_5$	$+t^4y_4 \wedge t^4y_5$	$-t^2y_5 \wedge t^4y_5$	$-t^3y_5 \wedge t^4y_5$
$-t^3y_2 \wedge t^5y_5$	$+t^2y_3 \wedge t^5y_5$	$-t^3y_3 \wedge t^5y_5$	$+t^3y_4 \wedge t^5y_5$	$-t^4y_4 \wedge t^5y_5$
$+t^3y_5 \wedge t^5y_5$	$-t^{-3}y_2 \wedge t^{-3}y_6$	$+t^{-2}y_2 \wedge t^{-3}y_6$	$-t^{-2}y_5 \wedge t^{-3}y_6$	$+t^{-1}y_5 \wedge t^{-3}y_6$
$+t^{-3}y_2 \wedge t^{-2}y_6$	$-t^{-1}y_2 \wedge t^{-2}y_6$	$+t^{-2}y_5 \wedge t^{-2}y_6$	$-y_5 \wedge t^{-2}y_6$	$+t^{-3}y_2 \wedge t^{-1}y_6$
$-t^{-2}y_2 \wedge t^{-1}y_6$	$+y_2 \wedge t^{-1}y_6$	$-t^{-4}y_3 \wedge t^{-1}y_6$	$+t^{-3}y_3 \wedge t^{-1}y_6$	$-t^{-1}y_3 \wedge t^{-1}y_6$
$-t^{-3}y_4 \wedge t^{-1}y_6$	$+t^{-2}y_4 \wedge t^{-1}y_6$	$-y_4 \wedge t^{-1}y_6$	$-t^{-3}y_5 \wedge t^{-1}y_6$	$+ty_5 \wedge t^{-1}y_6$
$+t^{-3}y_6 \wedge t^{-1}y_6$	$-t^{-2}y_6 \wedge t^{-1}y_6$	$-t^{-3}y_2 \wedge y_6$	$-t^{-2}y_2 \wedge y_6$	$+t^{-1}y_2 \wedge y_6$
$+2y_2 \wedge y_6$	$-2ty_2 \wedge y_6$	$+t^2y_2 \wedge y_6$	$+t^{-3}y_3 \wedge y_6$	$-t^{-2}y_3 \wedge y_6$
$-t^{-1}y_3 \wedge y_6$	$+3y_3 \wedge y_6$	$-ty_3 \wedge y_6$	$+t^2y_3 \wedge y_6$	$+t^{-2}y_4 \wedge y_6$
$-t^{-1}y_4 \wedge y_6$	$-y_4 \wedge y_6$	$+3ty_4 \wedge y_6$	$-t^2y_4 \wedge y_6$	$+t^3y_4 \wedge y_6$
$-y_5 \wedge y_6$	$+ty_5 \wedge y_6$	$-2t^2y_5 \wedge y_6$	$-t^{-2}y_6 \wedge y_6$	$+t^{-3}y_2 \wedge ty_6$
$+t^{-2}y_2 \wedge ty_6$	$-y_2 \wedge ty_6$	$-ty_2 \wedge ty_6$	$-t^3y_2 \wedge ty_6$	$-t^{-1}y_3 \wedge ty_6$
$+y_3 \wedge ty_6$	$-2ty_3 \wedge ty_6$	$-t^3y_3 \wedge ty_6$	$-y_4 \wedge ty_6$	$+ty_4 \wedge ty_6$
$-2t^2y_4 \wedge ty_6$	$-t^4y_4 \wedge ty_6$	$+t^{-2}y_5 \wedge ty_6$	$+t^{-1}y_5 \wedge ty_6$	$+t^2y_5 \wedge ty_6$
$+t^3y_5 \wedge ty_6$	$+t^{-1}y_6 \wedge ty_6$	$+2y_6 \wedge ty_6$	$-t^{-2}y_2 \wedge t^2y_6$	$-t^{-1}y_2 \wedge t^2y_6$
$+t^2y_2 \wedge t^2y_6$	$+t^{-1}y_3 \wedge t^2y_6$	$+t^2y_3 \wedge t^2y_6$	$+t^3y_3 \wedge t^2y_6$	$+y_4 \wedge t^2y_6$
$+t^3y_4 \wedge t^2y_6$	$+t^4y_4 \wedge t^2y_6$	$-t^{-1}y_5 \wedge t^2y_6$	$+ty_5 \wedge t^2y_6$	$-t^2y_5 \wedge t^2y_6$
$-t^4y_5 \wedge t^2y_6$	$-2y_6 \wedge t^2y_6$	$-ty_6 \wedge t^2y_6$	$+2y_2 \wedge t^3y_6$	$+t^4y_2 \wedge t^3y_6$

$$\begin{array}{ccccc}
-t^{-1}y_3 \wedge t^3y_6 & -t^3y_3 \wedge t^3y_6 & -y_4 \wedge t^3y_6 & -t^4y_4 \wedge t^3y_6 & -y_5 \wedge t^3y_6 \\
-t^2y_5 \wedge t^3y_6 & +t^5y_5 \wedge t^3y_6 & +y_6 \wedge t^3y_6 & +t^2y_6 \wedge t^3y_6 & -ty_2 \wedge t^4y_6 \\
-t^2y_2 \wedge t^4y_6 & -t^4y_2 \wedge t^4y_6 & +y_3 \wedge t^4y_6 & +ty_4 \wedge t^4y_6 & +ty_5 \wedge t^4y_6 \\
+t^2y_5 \wedge t^4y_6 & +t^4y_5 \wedge t^4y_6 & -t^5y_5 \wedge t^4y_6 & -ty_6 \wedge t^4y_6 & +t^3y_2 \wedge t^5y_6 \\
-t^2y_3 \wedge t^5y_6 & +t^3y_3 \wedge t^5y_6 & -t^3y_4 \wedge t^5y_6 & +t^4y_4 \wedge t^5y_6 & -t^3y_5 \wedge t^5y_6 \\
+t^3y_6 \wedge t^5y_6 & -t^4y_6 \wedge t^5y_6 & & & 
\end{array}$$

## References

- [Bi] J. Birman, *Braids, Links and Mapping Class Groups*, Ann. of Math. Stud., No. 82, Princeton Univ. Press, 1975.
- [Big] S. Bigelow, The Burau representation is not faithful for  $n = 5$ , *Geom. and Top.*, Vol. 3 (1999), pp.397–404.
- [CP] T. Church and A. Pixton, Separating twists and the Magnus representation of the Torelli group, arXiv:0804.3633, 2009.
- [Fox] R.H. Fox, Free differential calculus. I. Derivation in the free group ring, *Ann. of Math.* (2) 57 (1953), 547–560.
- [J1] D. Johnson, An abelian quotient of the mapping class group  $\mathcal{I}_g$ , *Math. Ann.* 249 (1980), no. 3, 225–242.
- [J2] D. Johnson, The structure of the Torelli group. I. A finite set of generators for  $\mathcal{I}$ , *Ann. of Math.* (3) 118 (1983) 423–442
- [LP] D.D. Long and M. Paton, The Burau representation is not faithful for  $n \geq 6$ , *Topology*, Vol. 32, No. 2, 439–447 (1993).
- [Ma] W. Magnus, On a theorem of Marshall Hall, *Ann. of Math.* (2) 40, (1939). 764–768.
- [Me] G. Mess, The Torelli groups for genus 2 and 3 surfaces, *Topology*, Vol. 31, No. 2, 775–790 (1992).
- [Mo] J.A. Moody, The Burau representation of the braid group  $B_n$  is unfaithful for large  $n$ , *Bull. Amer. Math. Soc.* 25 (1991), pp. 379–384.
- [Mor] S. Morita, Structure of the mapping class groups of surfaces: a survey and a prospect, in *Geometry and Topology Monographs*, Vol. 2, pp.349–406.
- [S1] M. Suzuki, The Magnus representation of the Torelli group  $\mathcal{I}_{g,1}$  is not faithful for  $g \geq 2$ , *Proc. Amer. Math. Soc.*, Vol. 130, No. 3, 909–914 (2001)
- [S2] M. Suzuki, On the kernel of the Magnus representation of the Torelli group, *Proc. Amer. Math. Soc.*, Vol. 133., No. 6, 1865–1872 (2005)

Dept. of Mathematics, University of Chicago  
5734 University Ave.  
Chicago, IL 60637  
E-mails: tchurch@math.uchicago.edu, farb@math.uchicago.edu